



Letters on Applied and Pure Mathematics

On nonlinear Sobolev equations with terminal observations in L^p spaces

Donal O'Regan ¹

1. School of Mathematical and Statistical Sciences, University of Galway, Ireland

Abstract

In this paper, we investigate the backward problem for the heat equation equipped with the time fractional conformable derivative. This problem is a generalization of the classical heat equation. We consider the problem with a nonlinear source function in a bounded domain. This problem is shown to be ill-posed, so we regularize the solution by the Fourier truncated method and we estimate the error term in the $L^p(\mathcal{D})$ space. An example to illustrate the theory is given in the final section.

Key words: Conformable derivative; Fourier truncation method; Regularization.

2020 Mathematics Subject Classification: 35A05; 35A08.

Article history: Received 05 March 2023; Revised 24 April 2023; Accepted 05 May 2023; Online 09 May 2023

1 Introduction

Various derivatives arise in problems in engineering, economics, physics, etc., for example in the study of the anomalous diffusion process in physics [1, 2], problems related to heat propagation [3, 4], problems in biological engineering [5, 6], problems with partial differential equations in image processing [7], signal processing [8], control theory [9, 10], and uncertainty models with random walks [11, 12]; see also [13–19, 23, 31, 33–37] and the references therein. The time fractional conformable derivative [23–25] with order $\alpha \in (0, 1)$, denoted by $\frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha}$, is defined as:

$$\frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} f(t) = \lim_{\delta \rightarrow 0} \frac{f(t + \delta t^{1-\alpha}) - f(t)}{\delta}, \quad (1.1)$$

for all $t > 0$. If we consider $(0, t_0)$, $t_0 > 0$ and $\lim_{t \rightarrow t_0^+} \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} f(t)$ exists, then $\frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} f(t_0) = \lim_{t \rightarrow t_0^+} \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} f(t)$.

This type of derivative has applications in quantum mechanics [26], the authors in [27] used the conformable Fourier series to express the solution of the conformable fractional heat equation, problems in Newtonian mechanics were investigated with a conformable derivative in [29], and in [28] the authors obtained the stochastic solution by running the processes corresponding to conformable Cauchy problems with nonlinear deterministic clocks; for more applications of the conformable derivative we refer the reader to [42]–[43].

There are also many papers on partial differential equations with different types of derivatives [20–22]. In this paper we will consider the conformable derivative and the heat equation, namely:

$$\begin{cases} \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} w(x, t) + (-\Delta)^s w(x, t) - b \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} \Delta w(x, t) = F(w(x, t)), & x \in \mathcal{D}, t \in (0, T), \\ w(x, t) = 0, & x \in \partial\mathcal{D}, t \in (0, T), \\ w(x, T) = g(x), & x \in \mathcal{D}. \end{cases} \quad (1.2)$$

Here $\mathcal{D} \subset \mathbb{R}^N$ ($N \geq 1$) is a bounded domain with the smooth boundary $\partial\mathcal{D}$, and $T > 0$.

In this work, we focus on studying the well-posedness and the regularization process. In Hadamard's sense, a problem is well-posed if it satisfies three conditions, where the third condition is "continuous dependence of the solution on the given data". For partial differential equation problems, this third condition is frequently not satisfied which results in an ill-posed problem. Therefore, in order for the obtained solution to depend continuously on the observed data, we need to modify the problem to a class of other well-posed problems. This technique is called the regularization method and we refer the reader to [32, 38–41] and the references therein. The regularization method mentioned above all estimate the convergence rate between the exact solution and the regularized solution based on the principle of choosing a priori or a posteriori parameter. To our best of our knowledge, most of the previous studies evaluated the error term mainly in the $L^2(\mathcal{D})$ space. In this article, we obtain results for regularized problems in $L^q(\mathcal{D})$ for $q \neq 2$. The evaluation in $L^q(\mathcal{D})$ spaces for $q \neq 2$ is quite difficult and to achieve this we use some embeddings between $L^q(\mathcal{D})$ and the Hilbert scales space $\mathbb{X}^r(\mathcal{D})$.

The paper is organized as follows. Section 2 gives some preliminaries to be used later. In section 3, we derive the formula of the mild solution to (1.2), and show that (1.2) is ill-posed in Hadamard's sense. Section 4 considers the error estimation and solution regularization. The last section presents some numerical examples to demonstrate the obtained results.

2 Preliminary results

Definition 2.1. Let us consider the operator $(-\Delta)^s$ on $\mathcal{H}_0^1(\Omega) \cap \mathcal{H}^2(\Omega)$, and assume that the operator $(-\Delta)^s$ has the eigenvalues λ_n such that $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \leq \dots$ which approach ∞ as n goes to ∞ , and $e_n(x)$ is the orthonormal basis in $L^2(\mathcal{D})$. For all $s \geq 0$, we define by $(-\Delta)^s$ the operator

$$(-\Delta)^s v := \sum_{n=1}^{\infty} \langle v, e_n \rangle \lambda_n^s e_n.$$

The Hilbert scale space is

$$\mathbb{X}^r(\mathcal{D}) = \left\{ f \in L^2(\mathcal{D}), \sum_{n=1}^{\infty} \lambda_n^{2r} \left(\int_{\mathcal{D}} f(x) e_n(x) dx \right)^2 < \infty \right\},$$

for any $r \geq 0$. Now $\mathbb{X}^r(\mathcal{D})$ is a Hilbert space with norm:

$$\|f\|_{\mathbb{X}^r(\mathcal{D})} = \left(\sum_{n=1}^{\infty} \lambda_n^{2r} \left(\int_{\mathcal{D}} f(x)e_n(x)dx \right)^2 \right)^{1/2}, \quad f \in \mathbb{X}^r(\mathcal{D}). \quad (2.1)$$

Lemma 2.2. (See [30]) *The following are true:*

$$\left. \begin{aligned} L^q(\mathcal{D}) &\hookrightarrow \mathbb{X}^\mu(\mathcal{D}), & \text{if } & -\frac{N}{4} < \mu \leq 0, & q &\geq \frac{2N}{N-4\mu}, \\ \mathbb{X}^\mu(\mathcal{D}) &\hookrightarrow L^q(\mathcal{D}), & \text{if } & 0 \leq \mu < \frac{N}{4}, & q &\leq \frac{2N}{N-4\mu}. \end{aligned} \right\} \quad (2.2)$$

3 Inverse initial problem (1.2)

The mild solution of (1.2) is obtained by the Fourier series $w(x, t) = \sum_{n=1}^{\infty} w_n(t)e_n(x)$, with

$w_n(t) = \int_{\mathcal{D}} w(x, t)e_n(x)dx$. We have

$$\begin{cases} \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} \langle w(\cdot, t), e_n \rangle + b\lambda_n \frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} \langle w(\cdot, t), e_n \rangle - \lambda_n^s \langle w(\cdot, t), e_n \rangle = \langle F(\cdot, t), e_n \rangle, & t \in (0, T), \\ \langle w(\cdot, 0), e_n \rangle = \langle w_0, e_n \rangle. \end{cases} \quad (3.1)$$

Thus

$$\frac{\mathcal{C}\partial^\alpha}{\partial t^\alpha} w_n(t) - \lambda_n^s (1 + b\lambda_n)^{-1} w_n(t) = (1 + b\lambda_n)^{-1} F_n(t), \quad \text{where } F_n(t) = \int_{\mathcal{D}} F(x, t)e_n(x)dx. \quad (3.2)$$

Based on [31], we have

$$\begin{aligned} \langle w(\cdot, t), e_n \rangle &= \exp(-\lambda_n^s (1 + b\lambda_n)^{-1} t^\alpha \alpha^{-1}) \left(\int_{\mathcal{D}} w_0(x)e_n(x)dx \right) \\ &+ (1 + b\lambda_n)^{-1} \int_0^t \mu^{\alpha-1} \exp\left(\lambda_n^s (1 + b\lambda_n)^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1}\right) F_n(\mu) d\mu. \end{aligned} \quad (3.3)$$

Letting $t = T$, it follows from (3.3) that

$$\begin{aligned} \left(\int_{\mathcal{D}} g(x)e_n(x)dx \right) &= \exp\left(-\lambda_n^s (1 + b\lambda_n)^{-1} T^\alpha \alpha^{-1}\right) \left(\int_{\mathcal{D}} w_0(x)e_n(x)dx \right) \\ &+ (1 + b\lambda_n)^{-1} \int_0^T \mu^{\alpha-1} \exp\left(\lambda_n^s (1 + b\lambda_n)^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1}\right) F_n(\mu) d\mu. \end{aligned} \quad (3.4)$$

From (3.4), we get

$$\begin{aligned} \left(\int_{\mathcal{D}} w_0(x)e_n(x)dx \right) &= \left(\exp\left(-\lambda_n^s (1 + b\lambda_n)^{-1} T^\alpha \alpha^{-1}\right) \right)^{-1} \left[\left(\int_{\mathcal{D}} g(x)e_n(x)dx \right) \right. \\ &\left. - (1 + b\lambda_n)^{-1} \int_0^T \mu^{\alpha-1} \exp\left(\lambda_n^s (1 + b\lambda_n)^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1}\right) F_n(\mu) d\mu \right]. \end{aligned} \quad (3.5)$$

Substituting (3.5) into (3.4) and we obtain

$$\begin{aligned} \langle w(\cdot, t), e_n \rangle &= \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g(x)e_n(x)dx\right) \\ &\quad - (1+b\lambda_n)^{-1} \int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(\mu) d\mu. \end{aligned} \quad (3.6)$$

This leads to

$$\begin{aligned} w(x, t) &= \sum_{n=1}^{+\infty} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g(x)e_n(x)dx\right) e_n(x) \\ &\quad - \sum_{n=1}^{+\infty} (1+b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(\mu) d\mu\right) e_n(x). \end{aligned} \quad (3.7)$$

From (3.7), we have

$$\begin{aligned} w(x, t) &= \sum_{n=1}^{+\infty} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g(x)e_n(x)dx\right) e_n(x) \\ &\quad - \sum_{n=1}^{+\infty} (1+b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(w(\mu)) d\mu\right) e_n(x). \end{aligned} \quad (3.8)$$

3.1 The ill-posedness of problem (1.2)

For any $k \in \mathbb{N}^*$, where \mathbb{N}^* is the set of non-negative natural numbers. Let g and $F(w)$ be

$$\begin{cases} g(x) := \lambda_k^{-1/2} e_k(x), \\ F(w) := \sum_{k=1}^{\infty} \frac{\alpha(1+b\lambda_k)}{2\mathcal{C}_F T^\alpha \exp\left(\frac{\lambda_k^s T^\alpha}{1+b\lambda_k \alpha}\right)} \left(\int_{\mathcal{D}} w(x)e_k(x)dx\right) e_k(x). \end{cases} \quad (3.9)$$

let w satisfy the integral equation

$$\begin{aligned} w(x, t) &= \sum_{n=1}^{+\infty} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g(x)e_n(x)dx\right) e_n(x) \\ &\quad - \sum_{n=1}^{+\infty} (1+b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(w(\mu)) d\mu\right) e_n(x). \end{aligned} \quad (3.10)$$

For $w \in L^\infty(0, T; L^2(\mathcal{D}))$, let

$$\begin{aligned} \mathcal{R}(w)(x, t) &= \sum_{n=1}^{+\infty} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g(x)e_n(x)dx\right) e_n(x) \\ &\quad - \sum_{n=1}^{+\infty} (1+b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(w(\mu)) d\mu\right) e_n(x). \end{aligned}$$

Then for any $w_1, w_2 \in L^\infty(0, T; L^2(\mathcal{D}))$, using Hölder’s inequality, we obtain

$$\begin{aligned} & \| \mathcal{R}(w_1)(\cdot, t) - \mathcal{R}(w_2)(\cdot, t) \|_{L^2(\mathcal{D})}^2 \\ &= \sum_{n=1}^{\infty} \frac{1}{(1 + b\lambda_n)^2} \left| \int_t^T \mu^{\alpha-1} \exp \left(\lambda_n^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \right. \\ &\quad \left. \times \left(\int_{\mathcal{D}} (F(w_1)(\mu) - F(w_2)(\mu)) e_n(x) dx \right) d\mu \right|^2 \\ &\leq \frac{\alpha}{4T^\alpha} \sum_{n=1}^{\infty} \left(\int_t^T \mu^{\alpha-1} d\mu \int_t^T \mu^{\alpha-1} \exp \left(2\lambda_n^{s-1} b^{-1} (\mu^\alpha - t^\alpha - T^\alpha) \alpha^{-1} \right) \right. \\ &\quad \left. \times \left(\int_{\mathcal{D}} ((w_1)(\mu) - (w_2)(\mu)) e_n(x) dx \right) d\mu \right) \\ &\leq \frac{\alpha}{4T^\alpha} \int_t^T \mu^{\alpha-1} \|w_1(\cdot, \mu) - w_2(\cdot, \mu)\|_{L^2(\mathcal{D})}^2 d\mu \leq 4^{-1} \|w_1 - w_2\|_{L^\infty(0, T; L^2(\mathcal{D}))}^2. \end{aligned} \tag{3.11}$$

This implies that

$$\| \mathcal{R}(w_1) - \mathcal{R}(w_2) \|_{L^\infty(0, T; L^2(\mathcal{D}))} \leq 2^{-1} \|w_1 - w_2\|_{L^\infty(0, T; L^2(\mathcal{D}))}.$$

Hence \mathcal{R} is a contraction. We conclude that $\mathcal{R}(w) = w$ has a unique solution $w \in L^\infty(0, T; L^2(\mathcal{D}))$ from Banach’s contraction theorem. From (3.10), we observe that

$$\begin{aligned} \|w(\cdot, t)\|_{L^2(\mathcal{D})} &\geq \left\| \underbrace{\sum_{n=1}^{\infty} \left(\exp \left(\lambda_n^s (1 + b\lambda_n)^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right) \int_{\mathcal{D}} g(x) e_n(x) dx \right) e_n(x)}_{\mathcal{R}_1(x, t)} \right\|_{L^2(\mathcal{D})} \\ &- \left\| \underbrace{\sum_{n=1}^{\infty} (1 + b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp \left(\lambda_n^s (1 + b\lambda_n)^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \left(\int_{\mathcal{D}} F(x, \mu, w(x, \mu)) e_n(x) dx \right) d\mu \right) e_n(x)}_{\mathcal{R}_2(x, t)} \right\|_{L^2(\mathcal{D})}. \end{aligned} \tag{3.12}$$

From $g(x) = \lambda_k^{-\frac{1}{2}} e_k(x)$ for any $k \in \mathbb{N}^*$ and note that $\|e_k\|_{L^2(\mathcal{D})} = 1$, we have

$$\begin{aligned} \| \mathcal{R}_1(\cdot, t) \|_{L^2(\mathcal{D})} &= \left(\sum_{n=1}^{\infty} \left| \exp \left(\lambda_n^s (1 + b\lambda_n)^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right) \left(\int_{\mathcal{D}} g(x) e_n(x) dx \right) \right|^2 \right)^{\frac{1}{2}} \\ &= \left(\sum_{n=1}^{\infty} \left| \exp \left(\lambda_n^s (1 + b\lambda_n)^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right) \left(\int_{\mathcal{D}} \lambda_k^{-\frac{1}{2}} e_k(x) e_n(x) dx \right) \right|^2 \right)^{\frac{1}{2}} \\ &\geq \lambda_k^{-\frac{1}{2}} \exp \left(\lambda_k^s (1 + b\lambda_k)^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right) \|e_k\|_{L^2(\mathcal{D})} \\ &= \lambda_k^{-\frac{1}{2}} \exp \left(\lambda_k^s (1 + b\lambda_k)^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right). \end{aligned} \tag{3.13}$$

By an argument similar to (3.11), we have

$$\| \mathcal{R}_2 u(\cdot, t) \|_{L^2(\mathcal{D})} \leq 2^{-1} \|u\|_{L^\infty(0, T; L^2(\mathcal{D}))}. \tag{3.14}$$

Combining (3.12)-(3.14), yields

$$\|u(\cdot, t)\|_{L^2(\mathcal{D})} \geq \lambda_k^{-\frac{1}{2}} \exp\left(\lambda_k^s(1+b\lambda_k)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) - \frac{1}{2} \|u\|_{L^\infty(0,T;L^2(\mathcal{D}))},$$

which implies that

$$\|u\|_{L^\infty(0,T;L^2(\mathcal{D}))} \geq \sup_{t \in [0,T]} \frac{2}{3} \lambda_k^{-\frac{1}{2}} \exp\left(\lambda_k^s(1+b\lambda_k)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right). \quad (3.15)$$

We also have

$$\sup_{t \in [0,T]} \exp\left(\lambda_k^s(1+b\lambda_k)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) = \exp\left(\lambda_k^{s-1}b^{-1}T^\alpha\alpha^{-1}\right). \quad (3.16)$$

Consequently, (3.15) becomes

$$\|u(\cdot, \cdot)\|_{L^\infty(0,T;L^2(\mathcal{D}))} \geq \frac{2}{3} \lambda_k^{-\frac{1}{2}} \exp\left(\lambda_k^{s-1}b^{-1}T^\alpha\alpha^{-1}\right). \quad (3.17)$$

Then, by letting $k \rightarrow \infty$, one has

$$\lim_{k \rightarrow \infty} \|g\|_{L^2(\mathcal{D})} = \lim_{k \rightarrow \infty} \lambda_k^{-\frac{1}{2}} = 0,$$

but

$$\lim_{k \rightarrow \infty} \|u\|_{L^\infty(0,T;L^2(\mathcal{D}))} \geq \frac{2}{3} \lim_{k \rightarrow \infty} \lambda_k^{-\frac{1}{2}} \exp\left(\lambda_k^{s-1}b^{-1}T^\alpha\alpha^{-1}\right) = \infty.$$

Therefore, problem (1.2) is ill-posed in the Hadamard sense in $L^\infty(0, T; L^2(\mathcal{D}))$.

4 Regularization of the nonlinear pseudo-parabolic equation on $L^q(\mathcal{D})$ spaces

For any $\mathcal{N}_{tr} > 0$, a regularized solution $\tilde{U}_\epsilon^{\mathcal{N}_{tr}}(x, t)$ is:

$$\begin{aligned} \tilde{U}_\epsilon^{\mathcal{N}_{tr}}(x, t) &= \sum_{n=1}^{\mathcal{N}_{tr}} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha - t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} g_\epsilon(x) e_n(x) dx\right) e_n(x) \\ &- \sum_{n=1}^{\mathcal{N}_{tr}} (1+b\lambda_n)^{-1} \left(\int_t^T \mu^{\alpha-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(\mu^\alpha - t^\alpha)\alpha^{-1}\right) F_n(\tilde{W}_\epsilon^{\mathcal{N}_{tr}}(\cdot, \mu)) d\mu\right) e_n(x). \end{aligned} \quad (4.1)$$

Theorem 4.1. Assume that $u \in L^\infty(0, T; \mathbb{X}^{a+\eta}(\mathcal{D}))$ for any $a > 0$ and $0 < \eta < \frac{N}{4}$, and there exists a constant $\mathcal{M} > 0$ such that

$$\mathcal{M} = \operatorname{ess\,sup}_{0 \leq t \leq T} \left(\sum_{n=1}^{\infty} \lambda_n^{2\eta_0} \exp\left(2t^\alpha\alpha^{-1}\lambda_n^{s-1}b^{-1}\right) |\langle u(t), e_n \rangle|^2 \right)^{\frac{1}{2}}, \quad (4.2)$$

where $\eta_0 > \eta$, the noisy data $g_\epsilon \in L^q(\mathcal{D})$ is such that

$$\|g_\epsilon - g\|_{L^q(\mathcal{D})} \leq \epsilon, \quad 1 < q < 2, \quad (4.3)$$

and \mathcal{N}_{tr} is chosen such that for any $t > 0$

$$\lim_{\epsilon \rightarrow 0} \mathcal{N}_{tr} = \infty, \quad \lim_{\epsilon \rightarrow 0} (\mathcal{N}_{tr})^\eta \exp\left(-\frac{Nq-2N}{4q} \epsilon\right) \exp\left((\mathcal{N}_{tr})^{s-1}b^{-1}T^\alpha\alpha^{-1}\right) \epsilon = 0. \quad (4.4)$$

Then $\|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^{\frac{2N}{N-4\eta}}(\mathcal{D})}$ is of order

$$\max \left\{ (\mathcal{N}_{tr})^{-a}, (\mathcal{N}_{tr})^{\eta-\eta_0}, (\mathcal{N}_{tr})^{\eta-\frac{Nq-2N}{4q}} \exp\left((\mathcal{N}_{tr})^{s-1}b^{-1}T^\alpha\alpha^{-1}\right) \epsilon \right\}. \quad (4.5)$$

Remark 4.2. By choosing $\mathcal{N}_{tr} = \left(\frac{b(1-\beta)}{\alpha^{-1}T^\alpha}\right)^{\frac{1}{s-1}} [\log(\epsilon^{-1})]^{\frac{1}{s-1}}$, for $\beta \in (0, 1)$, this implies that

$\|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^{\frac{2N}{N-4q}}(\mathcal{D})}$ is of order

$$\max \left\{ \left[\frac{b(1-\beta)}{\alpha^{-1}T^\alpha} \log(\epsilon^{-1}) \right]^{-\frac{\alpha}{s-1}}, \left[\frac{b(1-\beta)}{\alpha^{-1}T^\alpha} \log(\epsilon^{-1}) \right]^{\frac{\eta-\eta_0}{s-1}}, \left[\frac{b(1-\beta)}{\alpha^{-1}T^\alpha} \log(\epsilon^{-1}) \right]^{\eta - \frac{Nq-2N}{4q}} \epsilon^\beta \right\}.$$

Proof. The triangle inequality gives

$$\|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^2(\mathcal{D})} \leq \|\tilde{U}_\epsilon(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}}u(\cdot, t)\|_{L^2(\mathcal{D})} + \|u(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}}u(\cdot, t)\|_{L^2(\mathcal{D})}. \quad (4.6)$$

From (3.8), and let

$$\begin{aligned} \mathcal{P}_1(T-t)f &= \sum_{n=1}^{\infty} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha-t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} f(x)e_n(x)dx\right)e_n(x). \\ \mathcal{P}_2(T-t)f &= \sum_{n=1}^{\infty} (1+b\lambda_n)^{-1} \exp\left(\lambda_n^s(1+b\lambda_n)^{-1}(T^\alpha-t^\alpha)\alpha^{-1}\right) \left(\int_{\mathcal{D}} f(x)e_n(x)dx\right)e_n(x). \end{aligned} \quad (4.7)$$

We can rewrite (4.1) as:

$$\mathbb{T}_{\mathcal{N}_{tr}}u(t) = \mathcal{P}_1(T-t)\mathbb{T}_{\mathcal{N}_{tr}}g - \int_t^T \mu^{\alpha-1}\mathcal{P}_2(\mu-t)\mathbb{T}_{\mathcal{N}_{tr}}F(u(\mu))d\mu. \quad (4.8)$$

From (4.8), we have

$$\begin{aligned} \tilde{U}_\epsilon(t) - \mathbb{T}_{\mathcal{N}_{tr}}u(t) &= \mathcal{P}_1(T-t)\mathbb{T}_{\mathcal{N}_{tr}}(g_\epsilon - g) \\ &\quad - \int_t^T \mu^{\alpha-1}\mathcal{P}_2(\mu-t) \left[\mathbb{T}_{\mathcal{N}_{tr}}F(\tilde{U}_\epsilon(\mu)) - \mathbb{T}_{\mathcal{N}_{tr}}F(u(\mu))\right] d\mu. \end{aligned} \quad (4.9)$$

This implies that

$$\begin{aligned} \|\tilde{U}_\epsilon(t) - u(t)\|_{L^2(\mathcal{D})} &\leq \|\mathcal{P}_1(T-t)\mathbb{T}_{\mathcal{N}_{tr}}(g_\epsilon - g)\|_{L^2(\mathcal{D})} + \|u(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}}u(\cdot, t)\|_{L^2(\mathcal{D})} \\ &\quad + \left\| \int_t^T \mu^{\alpha-1}\mathcal{P}_2(\mu-t) \left[\mathbb{T}_{\mathcal{N}_{tr}}F(\tilde{U}_\epsilon(\mu)) - \mathbb{T}_{\mathcal{N}_{tr}}F(u(\mu))\right] d\mu \right\|_{L^2(\mathcal{D})} \\ &= \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3. \end{aligned} \quad (4.10)$$

For $1 < q < 2$, using the Sobolev embedding in Lemma 2.2, we get

$$\begin{aligned} \mathcal{I}_1 &= \|\mathcal{P}_1(T-t)\mathbb{T}_{\mathcal{N}_{tr}}(g_\epsilon - g)\|_{L^2(\mathcal{D})} \\ &\leq (\mathcal{N}_{tr})^{-\frac{Nq-2N}{4q}} \exp\left((\mathcal{N}_{tr})^{s-1}b^{-1}(T^\alpha-t^\alpha)\alpha^{-1}\right) \|g_\epsilon - g\|_{\mathbb{X}^{\frac{Nq-2N}{4q}}(\mathcal{D})} \\ &\leq \mathcal{C}_1(N, \eta)\mathcal{C}_2(N, q)(\mathcal{N}_{tr})^{-\frac{Nq-2N}{4q}} \exp\left((\mathcal{N}_{tr})^{s-1}b^{-1}(T^\alpha-t^\alpha)\alpha^{-1}\right) \|g_\epsilon - g\|_{L^q(\mathcal{D})}. \end{aligned} \quad (4.11)$$

Next,

$$\begin{aligned} \mathcal{I}_2 &= \left\| u(t) - \mathbb{T}_{\mathcal{N}_{tr}}u(t) \right\|_{L^2(\mathcal{D})} \\ &= \left(\sum_{\lambda_n > \mathcal{N}_{tr}} \exp\left(-2t^\alpha\alpha^{-1}\lambda_n^{s-1}b^{-1}\right) \lambda_n^{-2\eta_0} \exp\left(2t^\alpha\alpha^{-1}\lambda_n^{s-1}b^{-1}\right) \lambda_n^{2\eta_0} |\langle u(t), e_j \rangle|^2 \right)^{\frac{1}{2}} \\ &\leq (\mathcal{N}_{tr})^{-\eta_0} \exp\left(-t^\alpha\alpha^{-1}(\mathcal{N}_{tr})^{s-1}b^{-1}\right) \mathcal{M}, \end{aligned} \quad (4.12)$$

using $\lambda_n^{-2\eta_0} \leq (\mathcal{N}_{tr})^{-2\eta_0}$ for $\lambda_n > \mathcal{N}_{tr}$ and recalling

$$\mathcal{M} = \text{esssup}_{0 \leq t \leq T} \left(\sum_{n=1}^{\infty} \lambda_n^{2\eta_0} \exp \left(2t^\alpha \alpha^{-1} \lambda_n^{s-1} b^{-1} \right) |\langle u(t), e_n \rangle|^2 \right)^{\frac{1}{2}}. \quad (4.13)$$

A similar argument yields

$$\begin{aligned} \mathcal{I}_3 &= \left\| \mathcal{P}_2(\mu - t) \mathbb{T}_{\mathcal{N}_{tr}}(F(\tilde{U}_\epsilon(\mu)) - F(u(\mu))) \right\|_{\mathbb{X}^0(\mathcal{D})} \\ &\leq (1 + b\mathcal{N}_{tr})^{-1} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \|F(\tilde{U}_\epsilon(\mu)) - F(u(\mu))\|_{L^2(\mathcal{D})} \\ &\leq (1 + b\mathcal{N}_{tr})^{-1} L_F \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \|\tilde{U}_\epsilon(\mu) - u(\mu)\|_{L^2(\mathcal{D})} \\ &\leq (1 + b\mathcal{N}_{tr})^{-1} \mathcal{C}_3(N, \eta) L_F \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \|\tilde{U}_\epsilon(\mu) - u(\mu)\|_{L^2(\mathcal{D})}. \end{aligned} \quad (4.14)$$

We also find that

$$\begin{aligned} \mathcal{I}_3 &\leq (1 + b\mathcal{N}_{tr})^{-1} \mathcal{C}_3(N, \eta) L_F \\ &\quad \times \int_t^T \mu^{\alpha-1} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \|\tilde{U}_\epsilon(\mu) - u(\mu)\|_{L^2(\mathcal{D})} d\mu. \end{aligned} \quad (4.15)$$

Combining (4.10), (4.11), (4.12), and (4.15), we get

$$\begin{aligned} \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^2(\mathcal{D})} &\leq \|\tilde{U}_\epsilon(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}} u(\cdot, t)\|_{L^2(\mathcal{D})} + \|u(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}} u(\cdot, t)\|_{L^2(\mathcal{D})} \\ &\leq \mathcal{C}_1(N, \eta) \mathcal{C}_2(N, q) (\mathcal{N}_{tr})^{-\frac{Nq-2N}{4q}} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (T^\alpha - t^\alpha) \alpha^{-1} \right) \epsilon \\ &\quad + (\mathcal{N}_{tr})^{-\eta_0} \exp(-t^\alpha \alpha^{-1} (\mathcal{N}_{tr})^{s-1} b^{-1}) \mathcal{M} \\ &\quad + (1 + b\mathcal{N}_{tr})^{-1} \mathcal{C}_3(N, \eta) L_F \int_t^T \mu^{\alpha-1} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} (\mu^\alpha - t^\alpha) \alpha^{-1} \right) \|\tilde{U}_\epsilon(\mu) - u(\mu)\|_{L^2(\mathcal{D})} d\mu. \end{aligned}$$

From Gronwall's inequality, multiplying by $e^{\frac{t^\alpha}{\alpha} (\mathcal{N}_{tr})^{s-1} b^{-1}}$ on both sides, we get

$$\begin{aligned} e^{\frac{t^\alpha}{\alpha} (\mathcal{N}_{tr})^{s-1} b^{-1}} \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^2(\mathcal{D})} &\leq \mathcal{C}_1(N, \eta) \mathcal{C}_2(N, q) (\mathcal{N}_{tr})^{-\frac{Nq-2N}{4q}} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} T^\alpha \alpha^{-1} \right) \epsilon + (\mathcal{N}_{tr})^{-\eta_0} \mathcal{M} \\ &\quad + (1 + b\mathcal{N}_{tr})^{-1} \mathcal{C}_3(N, \eta) L_F \int_t^T \mu^{\alpha-1} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} \mu^\alpha \alpha^{-1} \right) \|\tilde{U}_\epsilon(\mu) - u(\mu)\|_{L^2(\mathcal{D})} d\mu. \end{aligned} \quad (4.16)$$

Thus

$$\begin{aligned} \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^2(\mathcal{D})} &\leq e^{-\frac{t^\alpha}{\alpha} (\mathcal{N}_{tr})^{s-1} b^{-1}} e^{(1+b\mathcal{N}_{tr})^{-1} (\mathcal{C}_3(N, \eta) L_F (T^\alpha - t^\alpha) \alpha^{-1})} \\ &\quad \times \left[\mathcal{C}_1(N, \eta) \mathcal{C}_2(N, q) (\mathcal{N}_{tr})^{-\frac{Nq-2N}{4q}} \exp \left((\mathcal{N}_{tr})^{s-1} b^{-1} T^\alpha \alpha^{-1} \right) \epsilon + (\mathcal{N}_{tr})^{-\eta_0} \mathcal{M} \right]. \end{aligned} \quad (4.17)$$

In view of the Sobolev embedding $\mathbb{X}^\eta(\mathcal{D}) \hookrightarrow L^{\frac{2N}{N-4\eta}}(\mathcal{D})$, we have that

$$\|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^{\frac{2N}{N-4\eta}}(\mathcal{D})} \leq \mathcal{C}_4(N, \eta) \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{\mathbb{X}^\eta(\mathcal{D})}. \quad (4.18)$$

Also

$$\|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{\mathbb{X}^\eta(\mathcal{D})} \leq \underbrace{\left\| \tilde{U}_\epsilon(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}} u(\cdot, t) \right\|_{\mathbb{X}^\eta(\mathcal{D})}}_{\mathcal{S}_1} + \underbrace{\|u(\cdot, t) - \mathbb{T}_{\mathcal{N}_{tr}} u(\cdot, t)\|_{\mathbb{X}^\eta(\mathcal{D})}}_{\mathcal{S}_2}. \quad (4.19)$$

Now, the first term \mathcal{S}_1 is estimated by

$$\mathcal{S}_1 \leq \left(\sum_{n=1}^{\lambda_n \leq \mathcal{N}_{tr}} \lambda_n^{2\eta} \left(\int_{\mathcal{D}} (\tilde{U}_\epsilon(x, t) - u(x, t)) e_n(x) dx \right)^2 \right)^{\frac{1}{2}} \leq (\mathcal{N}_{tr})^\eta \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^2(\mathcal{D})}. \quad (4.20)$$

Note $0 < \eta < \eta_0$ and \mathcal{S}_2 can be bounded as

$$\mathcal{S}_2 = \left(\sum_{\lambda_n > \mathcal{N}_{tr}} \lambda_n^{-2a} \lambda_n^{2a} \lambda_n^{2\eta} |\langle u(t), e_n \rangle|^2 \right)^{\frac{1}{2}} \leq (\mathcal{N}_{tr})^{-a} \|u\|_{L^\infty(0, T; \mathbb{X}^{a+\eta}(\mathcal{D}))}. \quad (4.21)$$

Combining (4.19) to (4.21), we conclude that

$$\begin{aligned} & \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{L^{\frac{2N}{N-4\eta}}(\mathcal{D})} \\ & \leq \mathcal{C}_4(N, \eta) \|\tilde{U}_\epsilon(\cdot, t) - u(\cdot, t)\|_{\mathbb{X}^\eta(\mathcal{D})} \leq \mathcal{C}_4(N, \eta) (\mathcal{N}_{tr})^{-a} \|u\|_{L^\infty(0, T; \mathbb{X}^{a+\eta}(\mathcal{D}))} \\ & \quad + \mathcal{C}_4(N, \eta) e^{-\frac{t^\alpha}{\alpha} (\mathcal{N}_{tr})^{s-1} b^{-1}} e^{(1+b\mathcal{N}_{tr})^{-1} (\mathcal{C}_3(N, \eta) L_F (T^\alpha - t^\alpha)^{\alpha-1})} \\ & \quad \times \left[\mathcal{C}_1(N, \eta) \mathcal{C}_2(N, q) (\mathcal{N}_{tr})^{\eta - \frac{Nq-2N}{4q}} \exp((\mathcal{N}_{tr})^{s-1} b^{-1} T^\alpha \alpha^{-1}) \epsilon + (\mathcal{N}_{tr})^{\eta - \eta_0} \mathcal{M} \right]. \end{aligned} \quad (4.22)$$

□

5 Simulation

In this section, we present some examples illustrating the theoretical part that we gave above. We recall the problem with some selected input data as follows

$$\frac{\mathcal{C} \partial^\alpha}{\partial t^\alpha} w(x, t) - \Delta w(x, t) - \frac{\mathcal{C} \partial^\alpha}{\partial t^\alpha} \Delta w(x, t) = F(x, t, w(x, t)), \quad (x, t) \in (0, \pi) \times (0, 1), \quad (5.1)$$

with the following Dirichlet boundary condition

$$w(0, t) = w(\pi, t) = 0, \quad t \in (0, 1), \quad (5.2)$$

adding the final value condition

$$w(x, 1) = g(x), \quad x \in (0, \pi), \quad (5.3)$$

where we choose the constants $T = 1, s = 1, b = 1$ and $\alpha \in (0, 1)$. Next, we set up some tools to support the calculation as follows.

Part 1. We approximate an integral by the composite Simpson rule as follows.

$$\int_{a_i}^{a_{i+1}} \mathcal{G}(s) ds \approx \frac{\Delta\tau}{3} \sum_{m=1}^{M/2} \left[\mathcal{G}(s_{2m-2}) + 4\mathcal{G}(s_{2m-1}) + \mathcal{G}(s_{2m}) \right], \quad (5.4)$$

where $s_m = a_i + m\Delta\tau$ for $m = 1, \dots, M-1, M$ and $\Delta\tau = \frac{a_{i+1} - a_i}{M}$.

Part 2. A uniform grid of mesh points (x_m, t_k) is employed to discretize the space and time intervals

$$x_m = \frac{(m-1)\pi}{M}, \quad m = 1, 2, \dots, M+1, \quad t_k = \frac{k-1}{N}, \quad k = 1, 2, \dots, N+1. \quad (5.5)$$

$\{t, \epsilon\}$	$\alpha = 0.1$		$\alpha = 0.5$		$\alpha = 0.9$	
	$RE_\alpha^\epsilon(t)$	$PE_\alpha^\epsilon(t)$	$RE_\alpha^\epsilon(t)$	$PE_\alpha^\epsilon(t)$	$RE_\alpha^\epsilon(t)$	$PE_\alpha^\epsilon(t)$
$t = 0.2, \epsilon = 1 \times 10^{-1}$	36.37	1.44 %	76.35	2.6 %	18.5	0.79 %
$t = 0.2, \epsilon = 3 \times 10^{-2}$	25.07	0.99 %	52.92	1.81 %	12.75	0.55 %
$t = 0.2, \epsilon = 5 \times 10^{-3}$	12.97	0.51 %	27.54	0.94 %	6.59	0.28 %

Table 1: The table error for $\alpha \in \{0.1, 0.5, 0.9\}$, $\epsilon \in \{1 \times 10^{-1}, 3 \times 10^{-2}, 5 \times 10^{-3}\}$ and $t = 0.2$.

The solution $u(x, t)$ can be presented in matrix form as follows

$$\begin{bmatrix} \mathcal{U}_{1,1} & \mathcal{U}_{1,2} & \cdots & \mathcal{U}_{1,N+1} \\ \mathcal{U}_{2,1} & \mathcal{U}_{2,2} & \cdots & \mathcal{U}_{2,N+1} \\ \vdots & \vdots & \vdots & \vdots \\ \mathcal{U}_{M+1,1} & \mathcal{U}_{M+1,2} & \cdots & \mathcal{U}_{M+1,N+1} \end{bmatrix}_{(M+1) \times (N+1)},$$

where the elements $\mathcal{U}_{m,n}$ are given by

$$\begin{aligned} \mathcal{U}_{m,n} &:= u(x_m, t_k) = \sqrt{\frac{2}{\pi}} \sum_{n=1}^{\mathcal{N}} U_n(t_k) \sin(nx_m) \\ &= \sqrt{\frac{2}{\pi}} \begin{bmatrix} U_1(t_k) & U_2(t_k) & \cdots & U_{\mathcal{N}}(t_k) \end{bmatrix} \times \begin{bmatrix} \sin(x_m) \\ \sin(2x_m) \\ \vdots \\ \sin(\mathcal{N}x_m) \end{bmatrix}, \end{aligned} \tag{5.6}$$

where \mathcal{N} is a parameter truncation,

$$\begin{aligned} U_n(t_k) &= \exp\left(\lambda_n^s (1 + b\lambda_n)^{-1} (1 - t_k^\alpha) \alpha^{-1}\right) \left(\int_0^\pi g(s) e_n(s) ds\right) \\ &- (1 + b\lambda_n)^{-1} \left(\int_{t_k}^1 \mu^{\alpha-1} \exp\left(\lambda_n^s (1 + b\lambda_n)^{-1} (\mu^\alpha - t_k^\alpha) \alpha^{-1}\right) \left(\int_0^\pi F(s, \mu, w(s, \mu)) e_n(s) ds\right) d\mu\right). \end{aligned} \tag{5.7}$$

Part 3. The relative error estimation RE_α^ϵ and percent error estimation are PE_α^ϵ

$$RE_\alpha^\epsilon(t) = \sum_{i=1}^{M+1} |u_\alpha(x_i, t) - u_\alpha^\epsilon(x_i, t)|, \quad PE_\alpha^\epsilon(t) = RE_\alpha^\epsilon(t) / \sum_{i=1}^{M+1} |u_\alpha(x_i, t)| \times 100. \tag{5.8}$$

In this example, we choose the input data

$$\begin{cases} F(x, t, w) = w^2 - [16(t^{2\alpha} + \alpha) + 34\alpha t^\alpha + (t^{4\alpha} + 2\alpha t^{2\alpha} + \alpha^2)] \sin(4x), & (x, t) \in (0, \pi) \times (0, 1), \\ g(x) = (1 + \alpha) \sin(4x), & x \in (0, \pi). \end{cases} \tag{5.9}$$

Let $\mathcal{N} = 10, M = N = 100$, and we have the following result

Tables 1, 2, 3 and Figure 1, we see that the numerical result is showing the convergent estimate of the relative error estimation RE_α^ϵ and percent error estimation PE_α^ϵ between the solutions. In this example, we consider three cases of ϵ such as $\epsilon \in \{1 \times 10^{-1}, 3 \times 10^{-2}, 5 \times 10^{-3}\}$ at $t \in \{0.2, 0.4, 0.6\}$. In Figure 1, we also show the 3D-graph of the solutions u on the domain $(x, t) \in [0, \pi] \times [0, 1]$.

$\{t, \varepsilon\}$	$\alpha = 0.1$		$\alpha = 0.5$		$\alpha = 0.9$	
	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$
$t = 0.4, \varepsilon = 1 \times 10^{-1}$	44.13	1.5 %	19.6	0.72 %	8.25	0.32 %
$t = 0.4, \varepsilon = 3 \times 10^{-2}$	30.64	1.04 %	13.59	0.50 %	5.77	0.22 %
$t = 0.4, \varepsilon = 5 \times 10^{-3}$	15.97	0.54 %	7.07	0.26 %	3.02	0.12 %

Table 2: The table error for $\alpha \in \{0.1, 0.5, 0.9\}$, $\varepsilon \in \{1 \times 10^{-1}, 3 \times 10^{-2}, 5 \times 10^{-3}\}$ and $t = 0.4$.

$\{t, \varepsilon\}$	$\alpha = 0.1$		$\alpha = 0.5$		$\alpha = 0.9$	
	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$	$RE_{\alpha}^{\varepsilon}(t)$	$PE_{\alpha}^{\varepsilon}(t)$
$t = 0.6, \varepsilon = 1 \times 10^{-1}$	9.65	0.35 %	2.1	0.08 %	2.39	0.09 %
$t = 0.6, \varepsilon = 3 \times 10^{-2}$	7.01	0.26 %	1.15	0.04 %	4.92	0.18 %
$t = 0.6, \varepsilon = 5 \times 10^{-3}$	3.81	0.14 %	0.44	0.02 %	2.39	0.09 %

Table 3: The table error for $\alpha \in \{0.1, 0.5, 0.9\}$, $\varepsilon \in \{1 \times 10^{-1}, 3 \times 10^{-2}, 5 \times 10^{-3}\}$ and $t = 0.6$.




Figure 1: An example figure

6 Conclusion

In this work, by applying the Fourier series truncation method, we introduced the regularized solution, and presented the error between the regularized solution and the exact solution derived in the space $L^q(\mathcal{D})$. We illustrated this with appropriate examples.

7 Declarations

Competing Interests

The author declare that they have no competing interests

Ethical Approval

Not applicable

Availability Data and Materials

Not applicable

Author's Contributions

All authors contributed equally. All the authors read and approved the final manuscript.

Funding

This research received no external funding.

References

- [1] K. Oldham, J. Spanier: *The Fractional Calculus Theory and Applications of Differentiation and Integration to Arbitrary Order*, Elsevier, (1974).
- [2] R. Metzler, J.H. Jeon, A.G. Cherstvy, E. Barkai, *Anomalous diffusion models and their properties: non-stationarity, non-ergodicity, and ageing at the centenary of single particle tracking*, Phys. Chem. Chem. Phys. **16**, pp. 24128–24164, 2014.
- [3] L.Dorcak and J. Terpak and I. Podlubny and L. PivkaL, *Methods for monitoring heat flow intensity in the blast furnace wall*, Metalurgija, **49** ,pp. 75–78, 2012.
- [4] J. Klafter and I. M. Sokolov: *Anomalous diffusion spreads its wings*, Phys World, **18**, pp. 29–32, 2005.
- [5] R.L. Magin: *Fractional Calculus in Bioengineering*, Begell House, Redding, (2006).
- [6] A. D. Freed and K. Diethelm: *Fractional calculus in biomechanics: a 3D viscoelastic model using regularized fractional derivative kernels with application to the human calcaneal fat pad*, Biomech. Model. Mechanobiol. **5**, pp. 203–215, 2006.
- [7] J. Bai, X. C. Feng: *Fractional-order anisotropic diffusion for image denoising*, IEEE. Trans. Image Process., **16** pp. 2492–2502, 2007.
- [8] R. J. I. I. Marks and M. Hall: *Differential integral interpolation from a bandlimited signal's samples*, IEEE. Trans. Acoust., **29**, pp. 872–877, 1981.
- [9] M. Caputo: *Linear models of dissipation whose Q is almost frequency independent—II*, Geophys. J. Int., **13** ,pp. 529–539, 1967;
- [10] I. Podlubny: *Fractional-order systems and fractional-order controllers*, Institute for Experimental Physics, Slovak Acad. Sci., **12**, 1–18, 1994.
- [11] R. Gorenflo and G. De Fabritiis and F. Mainardi: *Discrete random walk models for symmetric Levy–Feller diffusion processes*, Physica A. **269**, pp. 79–89, 1999.
- [12] R. Gorenflo and A. Vivoli: *Fully discrete random walks for space–time fractional diffusion equations*, Signal Process. **23**, pp. 2411–2420, 2003.
- [13] S. G. Samko and A.A. Kilbas and O.I. Marichev: *Fractional integrals and derivatives: Theory and applications*, Gordon and Breach Science Publishers, (1993).
- [14] A. A. Kilbas and H. M. Srivastava and J. J. Trujillo: *Theory and Applications of Fractional Differential Equations*, vol.204, North-Holland Mathematics Studies, Elsevier Science B.V., Amsterdam,(2006).
- [15] E. Karapinar, H.D. Binh, N.H. Luc, and N.H. Can: *On continuity of the fractional derivative of the time-fractional semilinear pseudo-parabolic systems*, Advances in Difference Equations, **1** 2021 , 1-24.
- [16] R. S. Adiguzel, U. Aksoy, E. Karapinar, I.M. Erhan: *On the solution of a boundary value problem associated with a fractional differential equation*, Mathematical Methods in the Applied Sciences. <https://doi.org/10.1002/mma.6652>
- [17] R. S. Adiguzel, U. Aksoy, E. Karapinar, I.M. Erhan: *Uniqueness of solution for higher-order nonlinear fractional differential equations with multi-point and integral boundary conditions*, RACSAM (2021) 115:155; <https://doi.org/10.1007/s13398-021-01095-3>

- [18] R. S. Adiguzel, U. Aksoy, E. Karapinar, I.M. Erhan: *On The Solutions Of Fractional Differential Equations Via Geraghty Type Hybrid Contractions*, Appl. Comput. Math., **20**, pp. 313-333, 2021.
- [19] N.D. Phuong, N.H. Luc, L.D. Long: *Modified Quasi Boundary Value method for inverse source problem of the bi-parabolic equation*, Advances in Theory of Nonlinear Analysis and its Applications, **4**, pp. 132–142, 2020.
- [20] M.A. Firoozjaee, S.A. Yousefi: *A numerical approach for fractional partial differential equations by using Ritz approximation*, Appl. Math. Comput., **338**, pp. 711-721, 2018.
- [21] T. A. Biala, A. Q. M. Khaliq: *Parallel algorithms for nonlinear time-space fractional parabolic PDEs*, J. Comput. Phys., **375**, pp. 135-154, 2018.
- [22] P. Prakash, S. Harikrishnan, M. Benchohra: *Oscillation of certain nonlinear fractional partial differential equation with damping term*, Appl. Math. Lett., **43** (2015), 72-79.
- [23] R. Khalil, M. Al Horani, A. Yousef, and M. Sababheh: *A new definition of fractional derivative*, Journal of Computational and Applied Mathematics, vol. **264**, pp. 65–70, 2014.
- [24] A. Atangana, D. Baleanu, and A. Alsaedi: *New properties of conformable derivative*, Open Mathematics, **13**, pp. 889–898, 2015.
- [25] T. Abdeljawad: *On conformable fractional calculus*, Journal of Computational and Applied Mathematics, **279**, pp. 57–66, 2015.
- [26] D. R. Anderson and D. J. Ulness: *Properties of the Katugampola fractional derivative with potential application in quantum mechanics*, Journal of Mathematical Physics, vol. **56**, (2015).
- [27] I. A. Hammad and R. Khalil: *Fractional Fourier series with applications*, American Journal of Computational and Applied Mathematics, vol. **4**, pp. 187–191, 2014.
- [28] Y. Çenesiz, A. Kurt, and E. Nane: *Stochastic solutions of conformable fractional Cauchy problems*, Statistics & Probability Letters, vol. **124**, 126–131, 2017.
- [29] W. S. Chung: *Fractional Newton mechanics with conformable fractional derivative*, Journal of Computational and Applied Mathematics, vol. **290**, 150–158, 2015.
- [30] N.H. Tuan, T. Caraballo: *On initial and terminal value problems for fractional nonclassical diffusion equations*, Proc. Amer. Math. Soc. **149**, no. 1, pp. 143–161, 2019.
- [31] N.H. Tuan, T.B. Ngoc, D. Baleanu, D. O'Regan: *On well-posedness of the sub-diffusion equation with conformable derivative model*, Communications in Nonlinear Science and Numerical Simulation, Volume **89**, (2020), 105332, 2020.
- [32] N.H. Tuan, L.D. Long, N.V. Thinh: *Regularized solution of an inverse source problem for a time fractional diffusion equation*, Applied Mathematical Modelling, Volume **40**, (2016) Issues 19–20, pp. 8244-8264, 2016.
- [33] N.H. Tuan, A.T. Nguyen, C. Yang: *Global well-posedness for fractional Sobolev-Galpern type equations.*, Discrete Contin. Dyn. Syst. **42**, no. 6, pp. 2637–2665, 2022.
- [34] N.H. Tuan, N.V. Tien, C. Yang: *On an initial boundary value problem for fractional pseudo-parabolic equation with conformable derivative*, Math. Biosci. Eng. **19**, no. 11, pp. 11232–11259, 2022.
- [35] N.H. Tuan, N.D. Phuong, T.N. Thach: *New well-posedness results for stochastic delay Rayleigh-Stokes equations*. Discrete Contin. Dyn. Syst. Ser. B, **28**, no. 1, 347–358, 2023.
- [36] de Andrade, Bruno; N. H. Tuan: *A non-autonomous damped wave equation with a nonlinear memory term*. Appl. Math. Optim., **85**, no. 3, Paper No. 23, 20 pp, 2022.
- [37] T. B. Ngoc, E. Nane, N.H. Tuan: *On a terminal value problem for stochastic space-time fractional wave equations*. Math. Methods Appl. Sci. **46**, no. 1, pp. 1206-1226, 2023.
- [38] A. Qian, Y. Li: *Optimal error bound and generalized Tikhonov regularization for identifying an unknown source in the heat equation*, J Math Chem ;**49**, 765–75, 2011.
- [39] S. Yang, X.T. Xiong, Y. Nie: *Iterated fractional Tikhonov regularization method for solving the spherically symmetric backward time-fractional diffusion equation*, Appl. Numer. Math. **160**, pp. 217–241, 2021.
- [40] F. Yang, J.L. Fu, P. Fan, X.X. Li: *Fractional Landweber iterative regularization method for identifying the unknown source of the time-fractional diffusion problem*, Acta Appl. Math., **175**, Paper No. 13, 19 pp, 2021.
- [41] F. Yang, P. Zhang. X.X. Li, *The truncation method for the Cauchy problem of the inhomogeneous Helmholtz equation*, Appl. Anal. **98**, no. 5, pp. 991–1004, 2019.
- [42] M. Bouaouid, K. Hilal, S. Melliani: *Nonlocal telegraph equation in frame of the conformable time-fractional derivative*, Adv. Math. Phys., Art. ID 7528937, 7 pp, 2019.
- [43] A.A. Omer, M. Maysaa, D. Baleanu, *New exact solution of generalized biological population model*, J. Nonlinear Sci. Appl. **10**, no. 7, pp. 3916–3929, 2017.